

NARCOTICS

WHAT ARE NARCOTICS?

Narcotics are drugs that relieve pain and often induce sleep. Some of these drugs are opiates, which include opium and its derivatives morphine, codeine, and heroin. Narcotics also include certain drugs that have a morphine-like action, such as methadone. Certain analogs of narcotic drugs have been developed, though they are not common drugs of abuse. For more information, see the Drug Analogs fact sheet in this series.

WHICH NARCOTICS ARE ABUSED?

All of them. Heroin accounts for most narcotic abuse in the United States. Medicinal narcotics are also abused, such as methadone, meperidine, morphine, and cough syrups and painkillers which contain codeine. Various narcotics may be sniffed, swallowed, or injected.

WHAT ARE THE RISKS OF NARCOTIC USE?

Physical dangers depend on the specific drug, its source, and the way it is used. Many problems are caused by uncertain dose levels, impure drugs, or mixing the narcotic with other drugs. Fatal overdose is possible. The risk caused by injecting the drug is much greater than taking the drug in other ways. An overdose can result in death minutes after the injection.

Infections from non-sterile solutions, syringes, and needles cause many diseases. These include HIV/ AIDS and serum hepatitis. Skin abscess, inflammation of the veins, and lung congestion also occur.

HOW MANY PEOPLE USE NARCOTICS?

According to the 1996 National Household Survey, heroin had been

used in the past year by 0.3 percent of those aged 12 to 17, by 0.9 percent of those aged 18 to 25, and 0.2 percent of those aged 26-34.

The national Monitoring the Future survey found in 1997 that use of heroin in the past month was reported by 0.6% of eighth graders, 0.6% of tenth graders, and 0.5% of twelfth graders. Use has increased since 1991.

In Michigan, the number of heroin users increased by nearly 75% from 1979 to 1995. In addition, use has spread since 1979 away from the city of Detroit and surrounding counties to smaller counties.

WHO BECOMES DEPENDENT ON HEROIN OR OTHER NARCOTICS?

Any person who takes anyone of these drugs can become dependent. Other factors may play a role, but use of the drug over time is the central reason. These drugs create a strong craving.

WHAT ARE THE WITHDRAWAL SYMPTOMS OF NARCOTICS?

When a dependent person stops taking the drug, withdrawal begins within 4-6 hours after the last use. Full symptoms begin 12-16 hours after the last dose. They may include shaking, sweating, vomiting, runny nose and eyes, muscle aches, chills, abdominal pains, and diarrhea. The intensity of any of these symptoms depends upon the degree of drug dependence.

WHAT IS THE TREATMENT FOR NARCOTIC DEPENDENCE?

Treatment most often includes medical care during withdrawal. Other drugs may be used to alleviate symptoms during withdrawal. Individual

and group counseling are part of most treatment programs. Support from a self help group such as Narcotics Anonymous is an important part of most recoveries. Help in learning new job and life skills is provided in many treatment programs.

Narcotic antagonists are drugs which block the "high" and other effects of narcotics. Research has found ways to use these drugs in treatment.

WHAT IS METHADONE?

Methadone is a synthetic narcotic painkiller. It is one of several drugs which may be used to treat dependence on heroin. It is given daily to relieve the physical craving for heroin and prevent onset of withdrawal symptoms. This allows the addict to lead a relatively normal life.

Since methadone itself causes dependence, its use is under strict regulation. Many programs later help the addict withdraw from methadone.

Fatal overdose from methadone can occur if people do not use the drug properly. Most methadone deaths occur when the drug is diverted to the street.

WHAT ARE THE RISKS OF NARCOTIC USE DURING PREGNANCY?

Narcotics used during pregnancy cause serious risks for the fetus. Fetal growth and brain development may be retarded. Fetal distress is more likely to occur. Injecting narcotics (or other drugs) also increases the risk that the baby will contract HIV/ AIDS or other diseases.

Narcotic use by a pregnant woman can lead to premature or stillborn babies. The infant can be born addicted and suffer severe withdrawal symptoms, including convulsions.

Babies born to heroin-addicted mothers are more likely to have serious and lasting medical problems. They are more likely to be hyperactive, have sleep problems, and show slowed muscle and speech development .

When supervised by a physician in a specialized women and pregnancy treatment program, methadone may be appropriate in treating a pregnant woman who is dependent on narcotics.

A woman who is pregnant, considering pregnancy, or breastfeeding should not use illegal narcotics. She should discuss with her physician any use of prescription narcotics and any other drugs.

LEGAL INFORMATION

Narcotics are Controlled Substances under Michigan and federal law. Use, possession, delivery, possession with intent to deliver, and manufacture of the drug are all illegal without appropriate license or prescription. Penalties include imprisonment and fines. Certain penalties are mandatory.

Penalties are increased if a person eighteen years or older distributes the drug to a person under eighteen, or distributes the drug near school property.

For details on the legal penalties, refer to the Michigan Law Fact Sheet in this series.

SOURCES

Johnston et al *Monitoring the Future Study*, 1997 data. Prev-Line website. University of Michigan press release December 20, 1997. Table 1b.

Johnston et al. *Monitoring the Future Study*, 1996 data. University of Michigan press release December 19, 1996. Table 1.

Michigan Department of Public Health/ Center for Substance Abuse Services. "Briefly... From CSAS" Summer 1995, Pages 2-3.

US Department of Health and Human Services/ ADAMHA. *Alcohol, Tobacco, and Other Drugs May Harm the Unborn*. 1990. Pages 35-40.

US Department of Health and Human Services/ Public Health Service. *National Household Survey on Drug Abuse: Population Estimates 1996*. 1997. Table 18, Page 103.

US Department of Labor. *What Works: Workplaces Without Drugs*. 1991, Page 62.

MRC Online!
Visit us today
www.WeAreMRC.org
Email: Info@WeAreMRC.org

MRC

MICHIGAN RESOURCE CENTER

For Alcohol, Tobacco & Other Drug Information,
Health Promotion, and Traffic Safety Educational Materials

111 WEST EDGEWOOD BLVD. STE 11
LANSING, MICHIGAN 48911
(517) 882-9955

MATERIALS INFO (800) 626-4636 • FAX (517) 882-7778 • E-mail: mrc@voyager.net

The Michigan Resource Center is operated by the Traffic Safety Association of Michigan under contracts from the Michigan Department of Community Health, Bureau of Substance Abuse Services and Health Promotion and Publication, including Substance Abuse Prevention and Treatment Block Grant Funds. Additional funding and materials are provided by the Michigan Office of Highway Safety Planning.